

Urine / Phlebotomy

10. Quality of venipuncture: Traumatic Clean



| | | | |
|---------------------|--------------------------------------|--|--|
| Mark all that apply | <input type="radio"/> Vein collapsed | <input type="radio"/> Excessive duration of draw | <input type="radio"/> Vein hard to get at |
| | <input type="radio"/> Hematoma | <input type="radio"/> Multiple sticks | <input type="radio"/> Leakage at venipuncture site |

If tube is not full, but is at least half full, please indicate "Partial" and enter the volume to the nearest mL.

| 11. Blood volume per tube: | Filled | | | Specify volume (mL): <i>min 1/2 full</i> |
|----------------------------|-----------------------|-----------------------|-----------------------|---|
| | Yes | No | Partial | |
| a. Serum 10 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| b. Serum 10 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| c. EDTA 10 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| d. EDTA 10 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| e. Paxgene 2.5 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| f. CBC/Diff EDTA 2 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |

For epigenetics sites:

| | | | | |
|-----------------------|-----------------------|-----------------------|-----------------------|----------------------|
| g. Cell Prep 8 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| h. Cell Prep 8 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| i. Cell Prep 8 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| j. Cell Prep 8 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| k. CBC/Diff EDTA 2 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |

For tissue sodium site:

| | | | | |
|------------------|-----------------------|-----------------------|-----------------------|----------------------|
| k. Heparin 10 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| l. Serum 5 mL | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |

(continued)

Urine / Phlebotomy

(continued)

11. Blood volume per tube:

| For PET site: | Filled | | | Specify volume (mL): <i>min 1/2 full</i> |
|----------------|-----------------------|-----------------------|-----------------------|--|
| | Yes | No | Partial | |
| m. 10 mL serum | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| n. 5 mL EDTA | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| o. 5 mL EDTA | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| p. 5 mL EDTA | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| q. 5 mL EDTA | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |
| r. 5 mL EDTA | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="text"/> |

12. Participant selected as a quality control subject?

(Participant ID ends in 6)

(Blood)

- NO
- YES
- YES, but not enough
blood for QC

(Urine)

- NO
- YES
- YES, but not enough
urine for QC

Comments: _____
