

(120-seconds optimum)



Biosample Collection

10. Time at evenipuncture		: O AM O PM	
11. Quality	of venipuncture: O	Fraumatic ○ Clean	
Mark all	O Vein collapsed	O Excessive duration of draw	O Vein hard to get at
that apply	O Hematoma	 Multiple sticks 	Leakage at venipuncture site

If tube is not full, but is at least half full, please indicate "Partial" and enter the volume to the nearest mL.

12. volume per tube:		Filled			Specify volume (mL): <i>min 1/2 full</i>
		Yes	No	Partial	
a.	EDTA 10 mL	0	0	0	
b.	EDTA 10 mL	0	0	0	
C.	EDTA 10 mL	0	0	0	
d.	EDTA 10 mL	0	0	0	
e.	Serum 10 mL	0	0	0	
f.	Serum 5 mL	0	0	0	
g.	EDTA 2 mL	0	0	0	
h.	EDTA 10 mL	0	0	0	
i.	PAXgene 2.5 mL	0	0	0	

(continued)



Biosample Collection

	C - I	
Hair	เดเ	lection

12	W/2c	hair	collection	comn	Chatal
13.	vvas	Hall	conection	COITID	ietea :

O Yes —

O No

13a. How many follicles were collected (1-10)?

13b. How many shafts were collected (0-10)?

13c. Location of hair collection:

O Front or head

O Left side

O Right side

О Тор

O Back

14. Is the participant selected as a quality control subject for serum, hair, and nasal brushing?

O NO

O YES -

Was serum QC sample collected?

Was hair QC sample collected?

Was nasal brushing QC sample collected?

0

0

NO

0

YES

0

Biosample QC ID:

First digit=FC, 2nd digit=9



Biosample Collection

15. Is the participant selected as a quality control subject for EDTA and urine?

O NO				
O YES		NO	YES	
	Was EDTA QC sample collected?	0	0	
	Was urine QC sample collected?	0	0	
			ļ	
	Bios	sample QC ID:		
	Firs	t digit=FC, 2nd o	digit=9	

Comments:			