



Exam 7 Medications Inventory

Craig Johnson

January 24, 2022

Importance of Medications

Exam 7 Ancillary Studies have a special interest in medications

- MESA-Mind Cognitive Assessment
 - memory, verbal and numeric reasoning, reaction time
 - Psychotropic drugs (amitriptyline, benzodiazepines, paracetamol, ...)
 - Sleep medications (ambien, night-time formulations, ...)
 - Calcium channel blockers
- MESA Sleep and 24 Hour Act
 - Sleep medications

Importance of Medications

- Of the 1,964 publications to date, >95% make use of medications data.

[Shah RV, Allison MA, Lima JA, Bluemke DA, Abbasi SA, Ouyang P, Jerosch-Herold M, Ding J, Budoff MJ, Murthy VL. Liver fat, statin use, and incident diabetes: The Multi-Ethnic Study of Atherosclerosis. *Atherosclerosis*. 2015;242\(1\):211-217.](#)

Table 1
Baseline characteristics of the study population, stratified by no statin at follow-up or use of statin at follow-up.

	All subjects (N = 3153)	No statin (N = 2237)	Incident statin use (N = 916)	P-value
Age (y)	59.0 [51.0–68.0]	58.0 [50.0–68.0]	61.0 [54.0–69.0]	<0.0001
Male gender	1384 (43.9)	992 (44.3)	392 (42.8)	0.43
Race				0.004
White	1235 (39.2)	834 (37.3)	401 (43.8)	
Asian	394 (12.5)	297 (13.3)	97 (10.6)	
Black	855 (27.1)	627 (28.0)	228 (24.9)	
Hispanic	669 (21.2)	479 (21.4)	190 (20.7)	
Weight (lb)	165.7 [143.0–192.0]	165.0 [142.3–190.0]	168.0 [144.1–194.9]	0.02
BMI (kg/m ²)	27.0 [24.1–30.3]	26.8 [23.9–30.2]	27.5 [24.6–30.9]	<0.0001
Waist circumference (cm)	95.0 [86.0–103.5]	94.2 [85.2–102.9]	96.5 [87.6–105.0]	<0.0001
Systolic BP (mmHg)	120.3 [108.0–136.0]	118.5 [107.0–134.5]	124.5 [112.5–140.0]	<0.0001
Diastolic BP (mmHg)	71.5 [64.5–78.0]	71.0 [64.5–77.5]	72.5 [65.5–79.0]	<0.0001
Hypertension	1116 (35.4)	693 (31.0)	423 (46.2)	<0.0001
Antihypertensive Rx	838 (26.6)	514 (23.0)	324 (35.4)	<0.0001
Smoking status				0.19
Never	1678 (53.2)	1181 (52.8)	497 (54.3)	
Former	1099 (34.9)	774 (34.6)	325 (35.5)	
Current	375 (11.9)	281 (12.6)	94 (10.3)	
Family history of diabetes	1062 (33.7)	726 (32.5)	336 (36.7)	0.02
Metabolic syndrome	644 (20.4)	390 (17.4)	254 (27.7)	<0.0001
Glucose (mg/dl)	86.0 [81.0–91.0]	86.0 [80.0–91.0]	87.0 [81.0–91.0]	0.002
CRP (mg/l)	1.7 [0.8–4.0]	1.6 [0.7–3.8]	2.0 [1.0–4.6]	<0.0001

Medications Form



Exam 7 Medications

Participant ID #:

Acrostic:

Technician ID:

Date: / /
Month Day Year

Section A Medication Reception

As you know, MESA will be describing all medications its participants are using, both **prescription** and **over-the-counter**. These include **pills, liquid medications, skin patches, eye drops, creams, salves, inhalers and injections**, as well as **cold or allergy medications, vitamins, herbal remedies and other supplements**. The letter you received about this appointment included a cloth medications bag for all your current medications and asked you to bring them to the clinic. Have you brought this bag with you? Are these all the medications that you have **taken in the past two weeks?**

- YES → May I see them? *Continue with Section B*
- NO → *Make arrangements to obtain*
- REFUSED → *Record reason for refusal in Comments Section*
- TOOK NO MEDICINES → *Go to end of form*



Medications Form

Section B Prescription Medications

1. Copy the name of the medicine, the strength (include units), and the total number of doses prescribed per day/week/month. Include all pills, skin patches, eye drops, creams, salves, injections, and inhalers (puffers).

2. On the average during the last two weeks, how many of these did you take a day/week/month?

Medication Name

Print the first 20 letters only - please print clearly

Strength (mg, IU, etc.)

Write the decimal as one of the digits

Number Prescribed

Circle: Day, Week, Month

__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M

PRN Medicine?

__	Y	N
__	Y	N
__	Y	N
__	Y	N
__	Y	N
__	Y	N
__	Y	N
__	Y	N
__	Y	N
__	Y	N

__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M



Medications Form

Section C Over-the-Counter Medications

3. Copy the name of the medicine, the strength (include units), and the total number of doses prescribed per day/week/month. Include all pills, liquid medications, eye drops, creams, salves, inhalers (puffers), and supplements.

Medication Name

Print the first 20 letters only - please print clearly

4. On the average during the last two weeks, how many of these did you take a day/week/month?

Strength (mg, IU, etc.)

Write the decimal as one of the digits

__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M
__	D	W	M




Practice Medications

	Medication	strength	Rx	Taken	Class	Comments
1	Losartan	25	1/day	1/day	Antihypertensive	Generic or trade OK
2	Atenolol	50	1/day	1/day	Beta blocker	Odd font
3	Lipitor	40	1/day	1 every other day	Statin	1/ 2 dys = 0.5 / day
4	Allopurinol	100	3/day	3 / 2 days	Gout Med	3 / 2 days = 1.5/day
5	Latanoprost	0.005%	2/day	2/day	Ocular	2doses (1 each eye)
6	Carvedilol	12.5	3/day	2/day	Beta blocker	2 nd BBlocker verify?
7	Ranitidine	150	2/day	1 every other day	H2 Blocker	PRN
8	Baclofen	20	qd	bid	Muscle relaxer	0.5 /day
9	Lantus	100/ML	As dir	15U/day	Insulin	Rx = -9 Taken = sum of typical dosing
10	Tylenol PM	500/25	OTC	qhs	NSAID	Anticholinergic
11	Aspirin	325	OTC	3/wk	NSAID	OTC PRN
12	Vitamin C	500	OTC	2/day	Supplement	Supplement

Losartan (antihypertensive)

Prescribed: 1/day

Taken: 1/day



Manuf. for: Merck Sharp & Dohme Corp., a subsidiary of
MERCK & CO., INC., Whitehouse Station, NJ 08889, USA
By: **MERCK SHARP & DOHME LTD.**
Cransington, Northumberland, UK NE23 3JU
Losartan Potassium (active ingred.) Made in France
Formulated in UK

NDC 0006-0951-54

Cozaar[®]
**(Losartan Potassium
Tablets)**

25 mg

Each tablet contains 25 mg of losartan potassium.
Store at 25°C (77°F); excursions permitted to 15-30°C
(59-86°F) [see USP Controlled Room Temperature].
Keep container tightly closed.
Protect from light.

Rx only

90 Tablets



7004027000

GTIN
00300060951541

USUAL DOSAGE: See Package Insert.

Atenolol (Beta Blocker)

Prescribed: 1/day

Taken: 1/day

USUAL DOSAGE: One tablet daily, or as directed by physician. Prescribing Information attached.

WARNING: As with all medications, keep out of the reach of children.

Store at controlled room temperature, 20–25°C (68–77°F) [see USP]. Dispense in well-closed, light-resistant containers.

35931-00

LOT

EXP

NDC 0310-0105-10 100 tablets

TENORMIN[®]
(atenolol)

50 mg tablets

Rx only

Manufactured for:
AstraZeneca Pharmaceuticals LP
Wilmington, DE 19850

By: IPR Pharmaceuticals, Inc.
Canóvanas, PR 00729 Product of Italy

AstraZeneca 

TENORMIN is a trademark of the AstraZeneca group of companies. © AstraZeneca 2012

3 N 0310-0105-10 2



Lipitor (Statin)

Prescribed: 1/day

Taken: 1 every other day

Store at controlled room temperature 20-25°C (68-77°F) [see USP].

Dispense in tight containers (USP).

DOSAGE AND USE
See package insert for full prescribing information.

*Each tablet contains atorvastatin calcium equivalent to 40 mg atorvastatin.

Distributed by
Parke-Davis
Division of Pfizer Inc
NY, NY 10017

MADE IN IRELAND

Pfizer

Pfizer

Pfizer

NDC 0071-0157-23

Lipitor[®]
(atorvastatin calcium)

40 mg*

tablets

90 Tablets Rx only

FPO: UPC @ 80%

N3 0071-0157-23 1

GTIN: 00300710157231
LOT:/EXP:

PAA056783

Allopurinol (Gout medication)

Prescribed: 3/day

Taken: 3 every other day

Made in USA

ZY001803

Rev. 10/03

Store at 15° to 25°C (59° to 77°F) in a dry place.
Dispense in light container as defined in the USP.
For indications, dosage, precautions, etc., see
accompanying package insert.

100 Tablets

NDC 65483-991-10

ZYLOPRIM®
(allopurinol)

Each scored tablet contains
100 mg

R_x Only


PROMETHEUS LABORATORIES INC.

Manufactured by DSM Pharmaceuticals, Inc., Greenville, NC 27834
for Prometheus Laboratories Inc., San Diego, CA 92121

October 2003

650392

LOT
EXP



Latanoprost (Ocular)

Prescribed: 2/day

Taken: 2/day

NDC 0517-0830-01

LATANOPROST

OPHTHALMIC SOLUTION 0.005%
(125 mcg/2.5 mL)

**FOR TOPICAL APPLICATION
IN THE EYE**

Sterile
Rx Only

AMERICAN
REGENT, INC.
SHIRLEY, NY 11967

Usual Dosage:

1 drop in the
affected eye(s) in
the evening.



RN1056-00
Rev. 12/10

LOT:
EXP:

Carvedilol (Beta Blocker)

Prescribed: 3/day

Taken: 2/day



A059855



4
0007-4141-20

Store below 30°C (86°F).
Dispense in a tight, light-resistant container.
Protect from moisture.
Each Tiltab® tablet contains carvedilol, 12.5 mg.
Dosage: See accompanying prescribing information.
Important: Use safety closures when dispensing this product unless otherwise directed by physician or requested by purchaser.

Manufactured for GlaxoSmithKline
Research Triangle Park, NC 27709
Manufactured by
Patheon Puerto Rico, Inc.
Manati, PR 00834, USA.
1000000058856 Rev: 8/08
Made in Ireland.

12.5mg
NDC 0007-4141-20

COREG®
CARVEDILOL TABLETS

100 TILTAB® Tablets



GlaxoSmithKline

R_x only

Ranitidine (H2 Blocker)

Prescribed: 2/day as needed

Taken: 1 every other day

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

Drug Facts

Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) **Purpose** Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor


Other information

- do not use if individual foil pouch is open or torn
- avoid excessive heat or humidity
- store at 20°-25°C (68°-77°F)
- this product is sodium and sugar free

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin


Questions? call 1-888-288-9199 (English/Spanish) M-F, 8:30-5 EST, or visit www.zantac.com for Tips for Managing Heartburn

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150



PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach


MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150



PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

TO OPEN
PUSH IN TAB AND PULL OUT

25 POUCHES
1 TABLET (1 DOSE)



What you should know about **MAXIMUM STRENGTH Zantac 150**
Ranitidine Tablets 150 mg/Acid Reducer

Please read all of this information before taking MAXIMUM STRENGTH Zantac 150.

What is MAXIMUM STRENGTH Zantac 150?

- MAXIMUM STRENGTH Zantac 150 contains 150 mg of ranitidine (as ranitidine hydrochloride, 168 mg), a medicine that doctors have prescribed more than 200 million times worldwide.

Excellent Safety Record

- The ingredient in MAXIMUM STRENGTH Zantac 150 ranitidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in MAXIMUM STRENGTH Zantac 150 has been taken safely with many frequently prescribed medications.
- MAXIMUM STRENGTH Zantac 150 tablets are sodium and sugar free.

What symptoms does MAXIMUM STRENGTH Zantac 150 relieve and prevent?

- MAXIMUM STRENGTH Zantac 150 relieves and prevents heartburn associated with acid indigestion and sour stomach.
- Certain foods or beverages, and even lying down to sleep can cause heartburn associated with acid indigestion and sour stomach. It is normal for the stomach to produce acid, especially after consuming food or beverages. However, acid in the wrong place, such as the esophagus, or too much acid, can cause burning pain and discomfort.

How should I take MAXIMUM STRENGTH Zantac 150?

- To relieve symptoms, swallow 1 tablet with a glass of water.
- To prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn.
- This medicine can be used up to twice daily (do not take more than 2 tablets in 24 hours).
- MAXIMUM STRENGTH Zantac 150 should not be given to children under 12 years old unless directed by a doctor.
- Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers.

How does MAXIMUM STRENGTH Zantac 150 work?


MAXIMUM STRENGTH Zantac 150 reduces the production of stomach acid. This is what makes MAXIMUM STRENGTH Zantac 150 different from antacids, which neutralize the acid already in your stomach. Antacids do not reduce the production of acid.

Tips for managing heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

When should I see a doctor?

- Do not use if you have trouble or pain swallowing food, vomiting with blood or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease except under the advice and supervision of a doctor.
- Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- Stop use and ask a doctor if your heartburn continues or worsens
- you need to take this product for more than 14 days
- If pregnant or breast-feeding, ask a healthcare professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
- Questions? call 1-888-288-9199 (English/Spanish) M-F, 8:30-5 EST, or visit www.zantac.com for Tips for Managing Heartburn.
- Do not use if individual foil pouch is open or torn.



Manufactured by: Boehringer Ingelheim (B) Consumer Health Care Products
Division of BI Pharmaceuticals, Inc., Ridgefield, CT 06877
© 2015, BI Pharmaceuticals, Inc. All rights reserved.
Product of Spain. Manufactured in Mexico.

Product repackaged and distributed by:
JC World Bell Wholesale Co., Inc.
376-380 Railroad Ave, Hackensack, NJ 07601

Do not discard this carton.
It contains important consumer information.


090320981/5
P011072-007
CT5971B
2/15 005



Montelukast (Leukotriene Inhibitor)

Prescribed: 1/day

Taken: 1/day



Singularair® 10 mg
(Montelukast Sodium) TABLETS
For Adults 15 Years of Age and Older

Dist. by:
MERCK & CO., INC.
Whitehouse Station, NJ 08889, USA

Manuf. by:
MERCK SHARP & DOHME LTD.
Cramlington, Northumberland, UK NE23 3JQ
Montelukast Sodium (active ingred.) Made in Ireland
Formulated in UK

Each tablet contains 10.4 mg
Montelukast Sodium equivalent to
10 mg Montelukast.

30 Tablets

Lot

NDC 0006-0117-31

Store at 25°C (77°F), excursions permitted to
15-30°C (59-86°F) [see USP Controlled Room
Temperature]. Protect from moisture and light.
Store in original package.

USUAL DOSAGE: See accompanying circular.

Rx only

9764000
30 | No. 3761

3 0006-0117-312

MARK
117

Baclofen (Muscle relaxer)

Prescribed: ½ tablet every day

Taken: ½ tablet / day

52959-733-02

RX Only: #XXXXXXXX

#XXX

CAUTION: Federal law PROHIBITS the transfer of this drug to anyone other than the person to whom prescribed and prohibits dispensing without a prescription unless OTC. See outsert for add'l RX info KEEP OUT OF REACH OF CHILDREN. Store in a cool dry place 68 to 77 degrees F.

BACLOFEN 10mg TABLET

Lot #: BCO114VX

Mfg: IVAX PHARM

Exp: 07/10


Mfg Croatia

Loc.:

Compare to: Lioresal

Mfg. NDC: 0172-4096-80

Pill ID: White Round Tablets



5295973302

Take as directed by your Doctor or
See outsert for usual dosage information



0172409680

BACLOFEN 10mg TABLET

52959-733-02

Qty #120

07/10

Lot BCO114VX

Lioresal

0172-4096-80

BACLOFEN 10mg TABLET

52959-733-02

Qty #120

07/10

Lot BCO114VX

Lioresal

0172-4096-80

BACLOFEN 10mg TABLET

52959-733-02

Qty #120

07/10

Lot BCO114VX

Lioresal

0172-4096-80

BACLOFEN 10mg TABLET

52959-733-02

Qty #120

07/10

Lot BCO114VX

Lioresal

0172-4096-80

Repack: HJ Harkins Co., Inc. Nipomo, CA 93444
Dispense in tight, child & light-resistant container per USP

Lantus (Insulin)

Prescribed: As directed

Taken: typically 5U in AM, 4U at midday, and 6U PM



Tylenol PM (NSAID)

Prescribed: Over the Counter (no prescription)

Taken: twice in the past 2 weeks



No. 26609 V1 Handy Solutions

Extra Strength
TYLENOL PM
2 POUCHES OF 2 CAPLETS EACH

Extra Strength
TYLENOL PM
2 POUCHES OF 2 CAPLETS EACH

Acetaminophen, Diphenhydramine HCl
Pain Reliever, Nighttime Sleep Aid Non-habit forming

2 POUCHES OF 2 CAPLETS EACH



DO NOT USE IF POUCH IS TORN OR DAMAGED

Drug Facts
Active ingredients (in each caplet)
Acetaminophen 500 mg Pain reliever
Diphenhydramine HCl 25 mg Nighttime sleep aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 4,000 mg of acetaminophen in 24 hours
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• with any other product containing diphenhydramine, even one used on skin
• in children under 12 years of age
• if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

Ask a doctor before use if you have
• liver disease
• a breathing problem such as emphysema or chronic bronchitis
• trouble urinating due to an enlarged prostate gland
• glaucoma

Ask a doctor or pharmacist before use if you are
• taking the blood thinning drug warfarin
• taking sedatives or tranquilizers

When using this product
• drowsiness will occur
• avoid alcoholic drinks
• do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if
• sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
• pain gets worse or lasts more than 10 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present • new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



LIFT HERE For More Drug Facts



Drug Facts (continued)
Directions
• do not take more than directed (see overdose warning)
adults and children 12 years and over • take 2 caplets at bedtime
• do not take more than 2 caplets of this product in 24 hours
children under 12 years do not use
Other information
• store between 20-25°C (68-77°F)
• do not use if pouch is torn or damaged
Inactive ingredients
carnauba wax, croscopolidone, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide
Questions or comments?
call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Handy Solutions® Guarantee If dissatisfied with this product, return it with a brief explanation to 5330 Fox Street, Denver, CO 80216. We'll replace it free of charge and refund your postage. You may also call 1-800-525-5097 for customer service, 9am to 4pm MST, weekdays or email us at cs@navajoinc.com. Visit us on the web at www.navajoinc.com

Repackaged and Distributed by: Handy Solutions® A Trademark of Navajo Mfg. Co. Inc. 5330 Fox Street Denver, CO 80216 Assembled in USA
Distributed By: McNeil Consumer Healthcare Division of McNEIL-PPC, Inc. Fort Washington, PA 19034 USA ©McNEIL-PPC, INC. 2015 Visit us at www.tylenol.com or call toll-free 1-877-TYLENOL (1-877-895-3665)

ASA (NSAID)

Prescribed: Over the counter (no prescription)

Taken: 3/week

GENUINE ASPIRIN
BAYER
ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER

THE WONDER DRUG

Safe Pain Relief Plus Lifesaving Benefits

325^{mg}

50 COATED TABLETS

To Open,
PUSH IN TAB and PULL OUT

Talk to your doctor or other healthcare provider before using this product for your heart.

25 Pouches of 2 Tablets Each

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
 - colds
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

(continued on back panel)

Drug Facts (continued)

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature

Inactive ingredients

corn starch, hypromellose, powdered cellulose, triacetin

Questions? 1-800-331-4536

Item #45650



3 47682 45650 4

Product of Spain

*This product is repackaged by:
Medique Products
17080 Alico Commerce Ct., Suite 1
Fort Myers, FL 33967
(1-800-634-7680
M-F 8:00am-4:00pm)
from a product Manufactured
by Bayer.

Bayer

Bayer, the Bayer Cross and
The Wonder Drug are registered
trademarks of Bayer, used with
permission.

Vitamin C (Supplement)

Prescribed: Over the counter (no prescription)

Taken: 2/day

